§ 180.364

Commodity	Parts per million
Cucumber	4.0
Eggplant	6.0
Fruit, citrus, group 10	20.0
Grape	5.0
Grape, raisin	20.0
Nut, tree, group 14	0.5
Papaya	2.0
Peach	10.0
Pear	15.0
Pistachio	0.5
Plum, prune, fresh	4.0
Plum, prune, dried	20.0
Strawberry	10.0
	1

(2) Tolerances are established for residues of the miticide/acaricide fenbutatin-oxide, including its metabolites and degradates, in or on the animal commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only the sum of fenbutatin-oxide, hexakis (2-methyl-2-phenylpropyl)

distannoxane, and its organotin metabolites, dihydroxybis(2-methyl-2-phenylpropyl) stannane and 2-methyl-2-phenylpropylstannoic acid, calculated as the stoichiometric equivalent of fenbutatin-oxide, in or on the commodity.

Commodity	Parts per million
Cattle, fat	0.5
Cattle, meat	0.5
Cattle, meat byproducts	0.5
Egg	0.1
Goat, fat	0.5
Goat, meat	0.5
Goat, meat byproducts	0.5
Hog, fat	0.5
Hog, meat	0.5
Hog, meat byproducts	0.5
Horse, fat	0.5
Horse, meat	0.5
Horse, meat byproducts	0.5
Milk, fat	0.1
Poultry, fat	0.1
Poultry, meat	0.1
Poultry, meat byproducts	0.1
Sheep, fat	0.5
Sheep, meat	0.5
Sheep, meat byproducts	0.5

(b) Section 18 emergency exemptions. [Reserved]

(c) Tolerances with regional registrations. A tolerance with regional registration, as defined in §180.1(1), is established for residues of the miticide/acaricide fenbutatin-oxide, including its metabolites and degradates, in or on the plant commodity in the table in this paragraph. Compliance with the

tolerance level specified in this paragraph is to be determined by measuring only fenbutatin-oxide, hexakis (2-methyl-2-phenylpropyl) distannoxane, in or on the commodity.

Commodity	Parts per million
Raspberry	10.0

(d) Indirect or inadvertent residues. [Reserved]

[65 FR 33713, May 24, 2000, as amended at 72 FR 41930, Aug. 1, 2007; 73 FR 5109, Jan. 29, 2008; 76 FR 23494, Apr. 27, 2011]

§180.364 Glyphosate; tolerances for residues.

(a) General. (1) Tolerances are established for residues of glyphosate, including its metabolites and degradates, in or on the commodities listed below resulting from the application of glyphosate, the isopropylamine salt of glyphosate, the ethanolamine salt of glyphosate, the dimethylamine salt of glyphosate, the ammonium salt of glyphosate, and the potassium salt of glyphosate. Compliance with the following tolerance levels is to be determined by measuring only glyphosate (N-(phosphonomethyl)glycine).

Commodity	Parts per million
Acerola	0.2
Alfalfa, seed	0.5
Almond, hulls	25
Aloe vera	0.5
Ambarella	0.2
Animal feed, nongrass, group 18	400
Artichoke, globe	0.2
Asparagus	0.5
Atemoya	0.2
Avocado	0.2
Bamboo, shoots	0.2
Banana	0.2
Barley, bran	30
Beet, sugar, dried pulp	25
Beet, sugar, roots	10
Beet, sugar, tops	10
Berry and small fruit, group 13-07	0.20
Betelnut	1.0
Biriba	0.2
Blimbe	0.2
Breadfruit	0.2
Cacao bean, bean	0.2
Cactus, fruit	0.5
Cactus, pads	0.5
Canistel	0.2
Canola, seed	20
Carrot	5.0
Chaya	1.0
Cherimoya	0.2
Citrus, dried pulp	1.5
Coconut	0.1
Coffee, bean, green	1.0

Environmental Protection Agency

Commodity	Parts per million
Corn, pop, grain	0.1
Corn, sweet, kernel plus cob with husk removed	3.5
Custord apple	210
Custard apple Date, dried fruit	0.2 0.2
Dokudami	2.0
Durian	0.2
Epazote	1.3
Feijoa	0.2 0.2
Fish	0.25
Fruit, citrus, group 10-10	0.50
Fruit, pome, group 11–10	0.20
Fruit, stone, group 12	0.2
Galangal, rootsGinger, white, flower	0.2 0.2
Gourd, buffalo, seed	0.1
Governor's plum	0.2
Gow kee, leaves	0.2
Grain, cereal, forage, fodder and straw, group 16, except field corn, forage and field corn,	100
stoverGrain, cereal, group 15 except field corn, pop-	100
corn, rice, sweet corn, and wild rice	30
Grass, forage, fodder and hay, group 17	300
Guava	0.2
Herbs subgroup 19A	0.2
Hop, dried cones	7.0 0.2
llamalmbe	0.2
Imbu	0.2
Jaboticaba	0.2
Jackfruit	0.2
Kava, roots Kenaf, forage	0.2 200
Leucaena, forage	200
Longan	0.2
Lychee	0.2
Mamey apple	0.2
Mango Mangosteen	0.2 0.2
Marmaladebox	0.2
Mioga, flower	0.2
Noni	0.20
Nut, pine	1.0
Nut, tree, group 14Oilseeds, group 20, except canola	1.0 40
Okra	0.5
Olive	0.2
Oregano, Mexican, leaves	2.0
Palm heart	0.2
Palm heart, leaves	0.2 0.1
Papaya	0.2
Papaya, mountain	0.2
Passionfruit	0.2
Pawpaw	0.2
Pea, dryPeanut	8.0 0.1
Peanut, hay	0.5
Pepper leaf, fresh leaves	0.2
Peppermint, tops	200
Perilla, tops	1.8
PersimmonPineapple	0.2 0.1
Pistachio	1.0
Pomegranate	0.2
Pulasan	0.2
Quinoa, grain	5.0
Rambutan	0.2 0.1
Rice, wild, grain	0.1
Rose apple	0.2

Commodity	Parts per million
Sapodilla	0.2
Sapote, black	0.2
Sapote, mamey	0.2
Sapote, white	0.2
Shellfish	3.0
Soursop	0.2
Spanish lime	0.2
Spearmint, tops	200
Spice subgroup 19B	7.0
Star apple	0.2
Starfruit	0.2
Stevia, dried leaves	1.0
Sugar apple	0.2
Sugarcane, cane	2.0
Sugarcane, molasses	30
Surinam cherry	0.2
Sweet potato	3.0
Tamarind	0.0
Tea, dried	1.0
Tea, instant	7.0
Teff, forage	100
Teff, grain	5.0
Teff, hay	100
Ti, leaves	0.2
	0.2
Ti, roots	0.2
Ugli fruit	
Vegetable, bulb, group 3–07Vegetable, cucurbit, group 9	0.20 0.5
Vegetable, foliage of legume, subgroup 7A, ex-	0.5
cept soybean	0.2
Vegetable, fruiting, group 8-10 (except okra)	0.10
Vegetable, leafy, brassica, group 5	0.2
Vegetable, leafy, except brassica, group 4	0.2
Vegetable, leaves of root and tuber, group 2,	
except sugar beet tops	0.2
Vegetable, legume, group 6 except soybean	
and dry pea	5.0
Vegetables, root and tuber, group 1, except car-	
rot, sweet potato, and sugar beet	0.20
Wasabi, roots	0.2
Water spinach, tops	0.2
Watercress, upland	0.2
Wax jambu	0.2
	0.2
Yacon, tuber	0.2

(2) Tolerances are established for residues of glyphosate, including its metabolites and degradates, in or on the commodities listed below resulting from the application of glyphosate, the isopropylamine salt of glyphosate, the ethanolamine salt of glyphosate, the dimethylamine salt of glyphosate, the ammonium salt of glyphosate, and the potassium salt of glyphosate. Compliance with the following tolerance levels is to be determined by measuring only glyphosate (phosphonomethyl)glycine) and its metabolite N-acetyl-glyphosate (N-acetyl-*N*-(phosphonomethyl)glycine; calculated as the stoichiometric equivalent of glyphosate).

Commodity	Parts per Million
Cattle, meat byproducts	5.0

§ 180.367

Commodity	Parts per Million
Corn, field, forage Corn, field, grain Corn, field, stover Egg Goat, meat byproducts Grain aspirated fractions Hog, meat byproducts Horse, meat byproducts Poultry, meat Poultry, meat byproducts Sheep, meat byproducts Soybean, forage Soybean, hay Soybean, hulls Soybean, seed	13 5.0 100 0.05 5.0 310.0 5.0 5.0 0.10 1.0 5.0 200.0

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[45 FR 64911, Oct. 1, 1980]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting §180.364, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and at www.fdsys.gov.

§ 180.367 n-Octyl bicycloheptenedicarboximide; tolerances for residues.

(a) General. A tolerance of 5 parts per million is established for residues of the insecticide synergist N-octyl bicycloheptene dicarboximide, including its metabolites and degradates, in or on all food items in food handling establishments where food and food products are held, processed, prepared and/or served, provided that the food is removed or covered prior to such use, except for bagged food in warehouse storage which need not be removed or covered prior to applications of formulations containing N-octyl bicycloheptene dicarboximide. Compliance with the tolerance level specified in this paragraph is to be determined only N-octyl measuring bicycloheptene dicarboximide, in or on the commodity.

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

 $[65~{\rm FR}~33713,~{\rm May}~24,~2000,~{\rm as}~{\rm amended}~{\rm at}~75~{\rm FR}~60243,~{\rm Sept.}~29,~2010]$

§ 180.368 Metolachlor; tolerances for residues.

(a) General. (1) Tolerances are established for the combined residues (free and bound) of the herbicide metolachlor, 2-chloro-N-(2- ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl)acetamide, and its metabolites, determined as the derivatives, 2- [(2-ethyl-6-methylphenyl)amino]-1-propanol and 4-(2-ethyl-6-methylphenyl)-2- hydroxy-5-methyl-3-morpholinone, each expressed as the parent compound in the following raw agricultural commodities:

Commodity	Parts per million
Almond, hulls	0.30
Animal feed, nongrass, group 18	1.0
Cattle, fat	0.02
Cattle, kidney	0.20
Cattle, liver	0.05
Cattle, meat	0.02
Cattle, meat byproducts, except kidney and liver	0.04
Corn, field, forage	6.0
Corn, field, grain	0.10
Corn, field, stover	6.0
Corn, pop, grain	0.10
Corn, pop, stover	6.0
Corn, sweet, forage	6.0
Corn, sweet, kernel plus cob with husks re-	
moved	0.10
Corn, sweet, stover	6.0
Cotton, gin byproducts	4.0
Cotton, undelinted seed	0.10
Dillweed	0.50
Egg	0.02
Goat, fat	0.02
Goat, kidney	0.20
Goat, liver	0.05 0.02
Goat, meat	0.02
Goat, meat byproducts, except kidney and liver Grass, forage	10
Grass, lorage	0.20
Horse, fat	0.20
Horse, kidney	0.02
Horse, liver	0.05
Horse, meat	0.02
Horse, meat byproducts, except kidney and liver	0.04
Milk	0.02
Nut, tree, group 14	0.10
Okra	0.50
Peanut	0.20
Peanut, hay	20
Peanut, meal	0.40
Potato	0.20
Poultry, fat	0.02
Poultry, meat	0.02
Poultry, meat byproducts	0.05
Safflower, seed	0.10
Sheep, fat	0.02
Sheep, kidney	0.20
Sheep, liver	0.05
Sheep, meat	0.02
Sheep, meat byproducts, except kidney and	
liver	0.04
Sorghum, grain, forage	1.0
Sorghum, grain, grain	0.30
Sorghum, grain, stover	4.0
Soybean, forage	5.0